

FAUNA AND ZOOGEOGRAPHY OF SPIDERS (ARANEAE) IN BULGARIA

Christo Deltshv: Institute of Zoology, Bulgarian Academy of Sciences, 1 Tsar
Osvoboditel Bld., 1000 Sofia, Bulgaria. E-mail: cdeltshv@zoology.bas.bg

ABSTRACT. Bulgaria is home to 975 species of spiders in 41 families. This number was established after a critical review of the existing literature and taxonomic review of the available collections. The spiders are distributed in all districts of Bulgaria, occurring in lowlands, forests, mountains, caves and urban territories. According to their current distribution the established 975 species can be split into 27 zoogeographical categories, grouped into five major chorotypes (Cosmopolitan, Holarctic, European, Mediterranean, Endemics). The largest number of species belongs to the widely distributed species in the Holarctic, but the most characteristic are the endemics. Their established number (76 species) is high and reflects the local character of the fauna. This phenomenon can be attributed to the relative isolation of the mountains compared with the lowlands in the context of paleo-environmental changes since the Pliocene.

Keywords: Europe, diversity, distribution, chorotypes

The first information on the spiders fauna of Bulgaria came from the end of 19th century (Pavesi 1876). Systematic investigation started in the beginning of 20th century by P. Drensky (1913, 1921, 1929, 1931, 1936a, b, 1937, 1938, 1939, 1940, 1942, 1943). Drensky (1936a) published the only catalogue of the spiders on the Balkan Peninsula in which 624 species from Bulgaria were reported. More recent publications are a result of intensive faunistic research after 1967 (Deltshv 1967, 1972a, b, 1973, 1974, 1977a, b, 1978, 1980, 1983a, b, c, 1984, 1985, 1987a, b, 1988, 1990, 1992, 1993, 1996, 1997a, b, 1998, 2003; Deltshv & Blagoev 1995, 1997, 2001; Helsingin et al. 1977 2001]; Blagoev & Deltshv 1989; Blagoev et al. 2002; Dimitrov 1993, 1994, 1996, 1997, 1999, 2003; Dimitrov & Lazarov 1999, 2002; Thaler et al. 1994; Lazarov 1998, 2003, 2004; Lazarov et al. 2001; Tzonev & Lazarov 2001). The accumulation of new data makes possible a critical taxonomic and faunistic review, together with a zoogeographic analysis.

METHODS

The material treated herein can be divided into two major parts: the first comprises a critical incorporation of all available literature records concerning the distribution of spiders in Bulgaria; the second concerns the original collections obtained from 1965–2002 during a field survey covering most of the districts in

Bulgaria, kept in the collections of Institute of Zoology, Bulgarian Academy of Sciences.

RESULTS AND DISCUSSION

The spider fauna is represented in Bulgaria by 975 species, included in 41 families and 285 genera. The number of species is high compared with the number of spiders recorded from other European countries with similar territories (Tables 1, 2). The number of families is also high compared with the data for the world: 110 (Platnick 2005; Austria 40, Germany 39, Switzerland 39 (Blick et al. 2002). Best represented are the families Linyphiidae (226 species or 23.2%), Gnaphosidae (98 species or 10%), Salticidae (91 species or 9.3%), Lycosidae (80 species or 8.2%) and Theridiidae (74 species or 7.5%). The genera with the highest number of species are: *Centromerus* (16 species or 7.3%), *Walckenaeria* (14 species or 6.4%), *Tenuiphantes* (11 species or 5%) and *Diplocephalus* (9 species or 4.1%) (Table 2). This richness, however, depends not only on the size of the regions, but also on the degree of exploration by araneologists.

According to their current distribution Bulgarian spiders can be divided into 27 zoogeographical chorotypes, grouped into 5 zoogeographical complexes (I = Cosmopolitan, II = Holarctic, III = European, IV—Mediterranean, V = Endemic) (Fig. 1). The data concerning general distribution of spiders are tak-

Table 1.—Comparison of area and spider species richness of some European countries.

Country	Area (km ²)	Spider species	Sources
Austria	83,858	961	Blick et al. (2002)
Bulgaria	110,993	975	Blagoev et al. (2002)
Czech Republic	77,280	830	Buchar & Růžička (2002)
Greece	128,900	810	Bosmans (pers. comm.)
Hungary	92,340	725	Samu & Szinetar (1999)
Macedonia	25,713	558	Blagoev (2002)
Portugal	91,500	660	Cardoso (1999)
Serbia	102,000	618	Deltshev et al. (2003)
Slovenia	20,120	529	Kuntner & Šereg (2002)

en from Michailov (1997), Marusik et al. (2000), Platnick (2004) and Vigna Taglianti et al. (1999) (Fig. 1).

Cosmopolitan species complex (COS + SCO, 20, 2%): Includes especially widespread species associated with lowlands, woodlands and high elevation zones of mountains.

Complex of species widely distributed in the Holarctic Region (HOL + OLW + PAT + PAL + WPA + ECA + EEC + SEC + EEE + WPA): Is best represented and comprises 561 (57.5%) species widespread in Bulgaria (Fig. 1). Palearctic species (sensu lato) are dominant (36.1%), followed by Holarctic (10.5%), European Central Asiatic (7%) and

West Palearctic (3%). The remaining chorotypes (EEC, SEC & EEE) are represented by single species. The complex includes especially widespread species associated with lowlands, woodlands and high elevation zones of mountains. Most of the species are well represented in the mountains. Characteristic mountain species are represented by the linyphiids *Bolyphantes alticeps* (Sundevall 1833), *B. luteolus* (Blackwall 1833), *Frontinellina frutetorum* (C. L. Koch 1834), *Goniatium rubens* (Blackwall 1833), *Pityohyphantes phrygianus* (C. L. Koch 1836), *Tenuiphantes alacris* (Blackwall 1853), *T. tenebricola* (Wider 1834). High mountain species are the lin-

Table 2.—The spider fauna of Bulgaria listed by families, depicting numbers of genera and species.

Families	Genera	Species	Families	Genera	Species
Atypidae	1	2	Oxyopidae	1	3
Nemesiidae	2	4	Zoropsidae	1	2
Filistatidae	2	2	Zoridae	1	6
Scytodidae	1	1	Agelenidae	6	32
Leptonetidae	1	2	Cybaeidae	2	3
Pholcidae	4	7	Hahnidae	4	9
Segestridae	1	3	Dictynidae	8	15
Dysderidae	4	29	Amaurobiidae	5	22
Oonopidae	4	4	Titanoecidae	2	7
Mimetidae	2	4	Miturgidae	1	12
Eresidae	1	2	Anyphenidae	1	2
Oecobiidae	1	1	Liocranidae	7	13
Uloboridae	2	4	Clubionidae	1	26
Nesticidae	1	3	Corinnidae	3	5
Theridiidae	17	74	Zodariidae	1	11
Theridiosomatidae	1	1	Gnaphosidae	19	96
Linyphiidae	94	226	Sparassidae	2	3
Tetragnathidae	4	17	Philodromidae	4	32
Araneidae	16	56	Thomisidae	13	60
Lycosidae	11	80	Salticidae	31	91
Pisauridae	2	3			
			Total	285	975




Figure 1.—Zoogeographical types in the spider fauna of Bulgaria, showing the number of species represented in each. Abbreviations: COS = cosmopolitan; SCO = subcosmopolitan; HOL = Holarctic; OLW = Old World; PAT = Palearctic-Paleotropical; PAL = Palearctic; WPA = west-Palearctic; ECA = European-central Asian; EEC = east European-central Asian; SEC = south European-central Asian; EEE = east European-east Mediterranean; MCA = Mediterranean-central Asian; BCA = Balkan-central Asian; EUR = European; MEE = middle-east European; MSEE = middle-southeast European; SEU = south European; EEU = east European; SEE = southeast European; BCAU = Balkan-Caucasian; BAN = Balkan Anatolian; MED = Mediterranean; EME = east Mediterranean; NME = north Mediterranean; NEM = northeast Mediterranean; BALK = Balkan endemics; BULG = Bulgarian endemics.

lyphiids *Entelecara media* (Kulczyński 1887) and *Mecynargus paetulus* (O.P.-Cambridge 1875), which are not established in the forest belt. Some xenotopic species (Thaler 1988) are widely distributed in the mountains and reach the highest summits as aeronauts. To this group belong the linyphiids *Dicymbium nigrum* (Blackwall 1834), *Diplostyla concolor* (Wider 1834), *Meioneta rurestris* (C.L. Koch 1836), *Oedothorax agrestis* (Blackwall 1853), *O. apicatus* (Blackwall 1850), *O. fuscus* (Blackwall 1834) which inhabit the mountain zone in dense populations (Deltshev 1990, 1995).

European species complex (EUR + MEE + MSEE + EEU + SEE): Comprises 191 (20%) species, widespread in Europe and Bulgaria (Fig. 1). European species (sensu lato) are dominant (14%), followed by East European species (3%), and Middle Southeast European species (1.5%). The remaining chorotypes (MEE & SEE) are represented by single species. The complex comprises widespread species which inhabit both lowland and mountains. Interesting is the group of European

mountain species, best represented in the forest, subalpine and alpine belts. Characteristic mountain species are the linyphiids, *Araeoncus anguineus* Deltshv 1987, *Bolyphantes kolosvaryi* (Caporiacco 1936), *Cinetata gradata* (Simon 1881), *Diplocephalus foraminifer* (O.P.-Cambridge 1875), *Improphantes improbulus* (Simon 1929), *Maso gallicus* Simon 1894, *Mughiphantes pulcher* (Kulczyński 1881), *Oreonetides glacialis* (C.L. Koch 1872), *Tiso vagans* (Blackwall 1834). Other linyphiid species such as *Palliduphantes istrianus* (Kulczyński 1914), *Centromerus capucinus* (Simon 1884), *C. cavernarum* (L. Koch 1872), *Porrhomma lativelum* Tretzel 1956 and *P. microps* (Roewer 1931), are characteristic of caves.

Mediterranean species complex (MCA—MED + EME + NME + NEM + SEU + BCA + BAN + BCAU): Includes 127 species (13%) that occur in the Mediterranean area or a part of it. The complex forms only 13% of the total spider fauna of Bulgaria, but the real percentage is probably higher, because a large part of the endemics have a Mediterranean or-

igin. Most of the species in the complex are widely distributed in the Mediterranean region. Very interesting are the mountain-Mediterranean species [*Aculepeira talishia* (Zavadsky 1902), *Pardosa incerta* 1905], which may be regarded as ancient elements in the high mountains.

Endemic species complex (BALK + BULG): Includes 76 species (10%) established in Bulgaria (35 species) and other territories of the Balkan Peninsula (41 species). The established number is high and reflects the local character of the fauna. The question about the status and distribution of endemic spiders found in Bulgaria is complicated. Some of them are found only in restricted areas, while others show wider distributions, sometimes even over the whole peninsula.

According to their origin, the endemics form two groups. Some of the species can be regarded as probable remnants of ancient Mediterranean mountain fauna (paleoendemics), and others came from the northern parts of Europe during the glacials and evolved under isolation on mountains during the interglacials (neoendemics). The curious is the distribution of the genus *Antrohyphantes* Dumitrescu 1971 (Linyphiidae), found only in the high elevation zone and in caves. It is related to the genus *Fageiella* Kratochvíl 1934 (Linyphiidae), an endemic from the caves of the western part of the Balkan Peninsula (Bosnia, Montenegro). Their allopatric distribution indicates that they had already separated before the establishment of the Vardar tectonic zones (Deltshev 1996). This suggests that these two genera are paleoendemics.

Concerning the formation of cave fauna, Deeleman-Reinhold (1976) wrote that "many European cave spiders are probably relics of populations of moist Tertiary forests". Due to the lack of knowledge, it is difficult to determine with certainty which of the cave spider endemics of Bulgaria are Tertiary and which are Quaternary elements. Nevertheless, the blind species in the family linyphiid, *Centromerus bulgarianus* (Drensky 1931), *Troglohyphantes drenskii* Deltshev 1973 and *Troglohyphantes bureschianus* Deltshev 1975, all species with primitive three branched paracymbia, also can be regarded as probable paleoendemics (Deltshev 1996).

The linyphiid spiders *Araeoncus clivifrons* Deltshev 1987, *Diplocephalus altimontanus*

Deltshev 1984, *Drepanotylus pirinicus* Deltshev 1992, *Erigone l. pirini* Deltshev 1983, *Incestophantes annulatus* (Kulczyński 1882), *Mughiphantes lithoclasticolus* Deltshev 1983, *Metopobactrus orbelicus* Deltshev 1985, known only from the high alpine parts of the Pirin and Rila Mountains are high alpine elements?. Here, also can be placed *Thenuiphantes drenskyi* Helsdingen 1977, occurring in the high elevation belts of Pirin, Rila, Central Stara Planina and Vitosha mountains. These species are regarded as derivative of their respective North or Middle European species (*Diplocephalus picinus* (Blackwall 1841), *Drepanotylus borealis* Holm 1945, *Erigone longipalpis* (Sundevall 1830), *Metopobactrus prominulus* (O.P.-Cambridge 1872), due to the disjunction of ranges during the glacial and interglacial (Deltshev 1996; Deltshev & Blagoev 1997). The largest fraction of endemics was encountered mainly in caves, coastal sites, woodlands and high altitude zones.

The presence of the 975 spider species shows that Bulgaria is a territory of considerable species richness. This conclusion is supported also by the existence of 76 endemic species. In a zoogeographical respect, the widely distributed spiders in the Holarctic region are dominant. However, the most characteristic faunal elements are the endemics. Their number is high, and their faunistic composition reflects the local character of the fauna. According to their origin the endemics belong to two principal faunistic complexes: Mediterranean and European. This phenomenon can be explained by the relative isolation of the mountains compared with the lowlands, in the context of palaeo-environmental changes that have occurred since the Pliocene.

ACKNOWLEDGMENTS

I am especially indebted to Dr. K. Thaler, Dr. J. Dunlop and E. Stojcoska for access to the materials in the collections of University of Innsbruck, Museum of Natural History, Berlin and Macedonian Museum of Natural History, and to my colleagues Gergin Blagoev, Stoyan Lazarov, Z. Hubenov and S. Abadjiev for discussion and helpful assistance.

LITERATURE CITED

Blagoev, G. 2002. Check list of Macedonian spiders (Araneae). *Acta Zoologica Bulgarica*. 54(3):9–34.

- Blagoev, G. & C. Deltshv. 1989. Biotopical distribution of wolf-spiders (Araneae, Lycosidae) in the Zemen Gorge, Southwestern Bulgaria. *Ecology, Bulgarian Academy of Sciences* 22:73–80.
- Blagoev, G., Deltshv, C. & S. Lazarov. 2002. The Spiders (Araneae) of Bulgaria. Institute of Zoology, Bulgarian Academy of Sciences. Online at <http://cl.bas.bg/bulgarianspiders/>
- Blick, T., A. Hänggi & K. Thaler. (2002). Checklist of the arachnids of Germany, Switzerland, Austria, Belgium and the Netherlands (Arachnida: Araneae, Opiliones, Pseudoscorpiones, Scorpiones, Palpigradi). Version 2002 June 1. Online at <http://www.AraGes.de/checkliste.html>
- Buchar, J. & V. Růžička. 2002. Catalogue of Spiders of the Czech Republic. Edited by P. Merrett. Praha: Peres. 351 pp.
- Cardoso, P. 1999. Portuguese spiders checklist. <http://www.geocites.com/RainForest/Vines/15197/checklist.html>
- Deeleman-Reinhold, C.L. 1976. Distribution patterns in European cave spiders. *Bulletin of the South African Speleological Association*: 25–35.
- Deltshv, C. 1967. On the studies of spiders (Araneae) in the Vitosha Mountain. *Bulletin de Institut de Zoologie et Musee, Sofia* 24:51–56.
- Deltshv, C. 1972a. A contribution to the study of spiders (Araneae) from the caves in Bulgaria. II. Genus *Lepthyphantes* in Bulgarian caves. *Bulletin de Institut de Zoologie et Musee, Sofia* 36: 137–147.
- Deltshv, C. 1972b. A new genus of Bulgarian cave spiders (*Protoleptoneta hulgarica*, n. g., n. sp., Leptonetidae). *International Journal of Speleology* 4:275–283
- Deltshv, C. 1973. A new *Troglohyphantes* from Bulgarian caves (Araneae, Linyphiidae). *International Journal of Speleology* 5:103–109.
- Deltshv, C. 1974. A new *Centromerus* from Bulgarian caves (Araneae, Linyphiidae). *International Journal of Speleology* 6:81–86.
- Deltshv, C. 1977a. A new *Protoleptoneta* from caves (Araneae, Leptonetidae). *Acta Zoologica Bulgarica* 1:3–8.
- Deltshv, C. 1977b. Genus *Nesticus* (Nesticidae, Araneae) from Bulgarian caves. In V. Panoš (ed.), *Proceedings of 6th International Congress of Speleology, Olomouc* 5:73–78.
- Deltshv, C. 1978. A new *Histopona* (Araneae, Agelenidae) from Bulgarian caves. *Acta Zoologica Bulgarica* 10:57–59.
- Deltshv, C. 1980. A contribution to the taxonomical study of *pallidus* group of genus *Lepthyphantes* Menge (Araneae, Linyphiidae) in Bulgaria. *Acta Zoologica Bulgarica* 16:44–56.
- Deltshv, C. 1983a. A contribution to the taxonomical study of *sylvaticus* group of genus *Centromerus* F. Dahl (Araneae, Linyphiidae) in Bulgaria. *Acta Zoologica Bulgarica* 21:53–58.
- Deltshv, C. 1983b. Notes on the spiders of genus *Erigone* Audouin (Araneae, Erigonidae) in Bulgaria. *Acta Zoologica Bulgarica* 22:71–75.
- Deltshv, C. 1983c. A contribution to the taxonomical and faunistical study of genus *Lepthyphantes* Menge (Araneae, Linyphiidae) from Pirin Mountain. *Acta Zoologica Bulgarica* 23:25–32.
- Deltshv, C. 1984. A new *Diplocephalus* species from Bulgarian mountains (Arachnida, Araneae, Erigonidae). *Reichenbachia* 22:91–93.
- Deltshv, C. 1985. A contribution to the study of the family Erigonidae (Araneae) from Pirin Mountain, Bulgaria, with a description of a new species (*Metopobactrus orbelicus* sp. n.). *Bulletin of the British Arachnological Society* 6:359–366.
- Deltshv, C. 1987a. A critical review of genus *Zodarion* Walckenaer (Araneae, Zodariidae) in Bulgaria. *Acta Zoologica Bulgarica* 33:19–25.
- Deltshv, C. 1987b. A critical review of genus *Araeoncus* Simon in Bulgaria, with description of a new species (*Araeoncus clivifrons* sp. n.) (Arachnida, Araneae, Erigonidae). *Reichenbachia* 25:97–102.
- Deltshv, C. 1988. The genus *Fageiella* Kratochvil and *Antrohyphantes* Dumitresco (Araneae, Linyphiidae, Lepthyphanteae) in the caves of Balkan Peninsula. *TUB-Documentation* 38:293–302.
- Deltshv, C. 1990. A critical review of genus *Coelotes* Blackwall in Bulgaria with description of a new species (*Coelotes drenskii* sp. n.) (Araneae, Agelenidae). *Acta Zoologica Bulgarica* 40:29–43.
- Deltshv, C. 1992. A critical review of family Theridiidae (Araneae) in Bulgaria. *Acta Zoologica Bulgarica* 43:13–22.
- Deltshv, C. 1993. The genus *Tegenaria* Latreille in Bulgaria: A critical review with description of two sibling species (Arachnida, Araneae: Agelenidae). *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck* 80:167–174.
- Deltshv, C. 1995. Spiders (Araneae) from the high altitude zone of Rila Mountain (Bulgaria). *Berichte Natwissenschaftlich-Medizinischen Verein Innsbruck* 82:217–225.
- Deltshv, C. 1996. The origin, formation and zoogeography of endemic spiders of Bulgaria (Araneae). *Revue Suisse de Zoologie hors serie* 141–151.
- Deltshv, C. 1997a. A new *Cybaeus* from the Mountains of Balkan Peninsula. *Reichenbachia* 32:1–4.
- Deltshv, C. 1997b. *Crypocina deelemanae* g. n., sp. n., a remarkable spider from the mountains of Montenegro (Yugoslavia) (Arachnida, Araneae, Hahnidae). *Revue Suisse de Zoologie* 104: 485–489.
- Deltshv, C. 1998. Spiders from the high altitude zone of Central Stara Planina Mountain (Bulgar-

- ia) (Araneae). *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck* 85: 213–221.
- Deltshev, C. 2003. A critical review of the spider species (Araneae) described by P. Drensky in the period 1915–1942 from the Balkan. *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck* 90:135–150.
- Deltshev, C. & G. Blagoev. 1995. A critical review of family Lycosidae (Araneae) in Bulgaria. *Revue Arachnologique* 10:171–198.
- Deltshev, C. & G. Blagoev. 1997. The spiders of Pirin Mountain (Bulgaria). Taxonomic, faunistic and zoogeographical analysis (Araneae). *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck* 84:269–286.
- Deltshev, C. & G. Blagoev. 2001. A critical check list of Bulgarian spiders (Araneae). *Bulletin of the British Arachnological Society* 12:110–138.
- Deltshev, C., B.P.M. Curcic & G. Blagoev. 2003. The Spiders of Serbia. Institute of Zoology, Faculty of Biology, University of Belgrade. 833 pp.
- Dimitrov, D. 1993. New and rare spiders (Araneae) to the fauna of Bulgaria. Second National Scientific Entomological Conference Sofia. Pp. 73–75.
- Dimitrov, D. 1994. A record of *Achaearanea tabulata* from the Balkan Peninsula (Araneae: Theridiidae). *Arachnologische Mitteilungen* 8: 77–79.
- Dimitrov, D. 1996. *Coelotes deltshevi* sp. n., a new spider species from Bulgaria (Arachnida, Araneae, Zodariidae). *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck* 83:159–161.
- Dimitrov, D. 1997. Description of *Harpactea strandjica*, sp. n. from Bulgaria (Araneae, Dysderidae). *Bulletin of the British Arachnological Society* 10:322.
- Dimitrov, D. 1999. The spider fauna of the Strandzha Mountain (South-East Bulgaria). I. Faunistic data and taxonomic remarks (Arachnida: Araneae). *Acta Zoologica Bulgarica* 52:15–26.
- Dimitrov, D. 2003. *Erigonoplus spinifemoralis* sp. n. (Araneae: Linyphiidae)—a new species from Bulgaria. *Acta Zoologica Bulgarica* 55:33–35.
- Dimitrov, D. & S. Lazarov. 1999. Two new species of *Harpactea* from Bulgaria (Araneae: Dysderidae). *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck* 86:127–129.
- Dimitrov, D. & S. Lazarov. 2002. A contribution to the study of the spiders (Araneae) in Chepun Mountain and Dragoman Swampland (NW Bulgaria). *Acta Zoologica Bulgarica* 54(2):47–53.
- Drensky, P. 1913. Über die Spinnenfauna Bulgariens. *Spisanie Bulgarska Akadademia na Naukata* 2:1–146.
- Drensky, P. 1921. Contribution à l'étude des araignées de la Macédoine orientale et de Pirine planina. *Spisanie Bulgarska Akadademia na Naukata* 23:1–50.
- Drensky, P. 1929. Spinnen (Araneae) aus Mittel und Süd-West Mazedonien. *Spisanie Bulgarska Akadademia na Naukata* 39:1–76.
- Drensky, P. 1931. Höhlen-Spinnen aus Bulgarien. *Spisanie Bulgarska Akadademia Na Naukata* 44: 1–50.
- Drensky, P. 1936a. Katalog der echten Spinnen (Araneae) der Balkanhalbinsel. *Sbornik Bulgarskata Akademia na Naukata* 32:1–223.
- Drensky, P. 1936b. Studien über die Bulgarischen Spinnenfauna ihre ökologischen und biogeographischen Besonderheiten. *Travaux Societe Bulgare des Sciences Naturelles* 17:71–115.
- Drensky, P. 1937. Die Spinnenfauna Bulgariens. I. Unterordnung Mygalomorphae, Family Ctenizidae und Atypidae. *Izvestia Tsarski Prirodonauchni Instituti* 10:259–280.
- Drensky, P. 1938. Die Spinnenfauna Bulgariens. II. Unterordnung Arachnomorphae, I Gruppe Tetrastica, Familien: Filistatidae, Dysderidae und Oonopidae. *Izvestia Tsarski Prirodonauchni Instituti* 11:81–113.
- Drensky, P. 1939. Die Spinnenfauna Bulgariens. III. Unterordnung Arachnomorphae, II Gruppe Trionichia, Familien: Urocteidae, Uloboridae, Sicaridae, Pholcidae, Eresidae. *Izvestia Tsarski Prirodonauchni Instituti* 12:231–252.
- Drensky, P. 1940. Die Spinnenfauna Bulgariens. IV. Unterordnung Arachnomorphae, II Gruppe Trionichia, Familien: Zodariidae, Dictynidae und Amaurobiidae. *Izvestia Tsarski Prirodonauchni Instituti* 13:169–194.
- Drensky, P. 1942. Die Spinnenfauna Bulgariens. V. Unterordnung Arachnomorphae, II Gruppe Trionichia, Familie: Agelenidae. *Izvestia Tsarski Prirodonauchni Instituti* 15:33–60.
- Drensky, P. 1943. Die Spinnenfauna Bulgariens. VI. Unterordnung Arachnomorphae, II Gruppe Trionichia, Familie Euetrioidae. *Izvestia Tsarski Prirodonauchni Instituti* 16:219–254.
- Helsdingen, van, P.I., K. Thaler & C. Deltshev. 1977. The *tenuis* group of *Lepthyphantes* Menge (Araneae, Linyphiidae). *Tijdschrift voor Entomologie* 120:1–54.
- Helsdingen, van, P.I., K. Thaler & C. Deltshev. 2001. The European species of *Bolyphantes* with an attempt of a phylogenetic analysis (Araneae, Linyphiidae). *Memorie della Societa Entomologica Italiana* 80:3–35.
- Kuntner, M. & I. Šereg. 2002. Additions to the spider fauna of Slovenia, with a comparison of spider species richness among European countries. *Bulletin of the British Arachnological Society* 12:185–195.
- Lazarov, S. 1998. A contribution to the study of the spiders (Araneae) in Sushtinska Sredna Gora

- Mountain, Bulgaria. *Historia Naturalis Bulgaria* 9:27–34.
- Lazarov, S. 2003. A review of the family Dysderidae (Araneae) in Bulgaria: faunistic and zoogeographical analysis. Pp. 259–265. *In* *European Arachnology 2002* (F. Samu & C. Szinetár, eds) Plant Protection Institute and Berzsenyi College, Budapest
- Lazarov, S. 2004. A contribution to the study of spiders (Araneae) in Macedonia. *Acta Zoologica Bulgaria* 56:155–166.
- Lazarov, S., C. Deltchev & G. Blagoev. 2001. Spiders (Araneae) of Sushtinska Sredna Gora Mountain, Bulgaria. *Acta Zoologica Bulgaria* 53:3–28.
- Marusik, Y.M., D.V. Logunov & S. Koponen. 2000. Spiders of Tuva, South Siberia. Magadan. IBPN FEB RAS, 2000. 252 pp.
- Michailov, K.G. 1997. Catalogue of the spiders of the territories of the former Soviet Union (Arachnida, Aranei). Moscow, Zoological Museum of the Moscow State University. 416 pp.
- Pavesi P. 1876. Gli Arachnidi Turchi. *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale* 19:1–27.
- Platnick, N. 2004. *The World Spider Catalog, Version 4.5* American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog81-87/index.html>.
- Samu, F. & C. Szinetár. 1999. Bibliographic checklist of the Hungarian spider fauna. *Bulletin of the British Arachnological Society* 11:161–184.
- Thaler, K. 1988. Arealformen in der nivalen Spinnenfauna der Ostalpen (Arachnida, Aranei). *Zoologischer Anzeiger* 220:233–244.
- Thaler, K., P. Van Helsing & C. Deltchev. 1994. Vikariante Verbreitung im Artenkomplex von *Lepthyphantes annulatus* in Europa und ihre Deutung (Araneae, Linyphiidae). *Zoologischer Anzeiger* 232:111–127.
- Tzonev, G. & S. Lazarov. 2001. A contribution to the study of spiders (Araneae) in Osogovo Mountain, south-west Bulgaria. *Acta Zoologica Bulgaria* 53:67–78.
- Vigna Taglianti, A., P.A. Audisio, M. Biondi, M.A. Bologna, G.M. Carpaneto, A. De Biase, S. Fattorini, E. Piattella, R. Sindaco, A. Venchi & M. Zapparoli. 1999. A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. *Biogeographia* 20:31–59.

Manuscript received 10 June 2005, revised 15 September 2005.